
LECTURE NOTES
DISCRETE MATHEMATICS

CHAPTER 8

Modular arithmetic



Updated: 19 November 2026

8 Modular arithmetic

Alice, Bob, Charlie and David take turns checking a shared printer, in that order, then the cycle repeats. Number their positions $0, 1, 2, 3$. Starting with Alice at turn 0, who checks at turn 17? Listing all the turns would work, but four complete cycles use sixteen turns and return to Alice. One further turn reaches Bob. The remainder after division by four identifies the position in the cycle.

Chapter 3 used integer multiples in proofs; Chapter 7 grouped integers according to an even difference. We now develop remainders systematically and group integers by the same remainder. These groups support arithmetic on a finite set of labels, even though the integers themselves extend without bound.

What you should be able to do

After working through this chapter, you should be able to:

- justify divisibility claims using integer witnesses;
- find a quotient and remainder, including for negative integers;
- interpret congruence using remainders and divisibility;
- describe residue classes and their partition of the integers;
- calculate with congruences and diagnose invalid cancellation;
- specify modular addition and multiplication as binary operations;
- check closure and identify two-sided identities in small systems.

8.1 Divisibility

We work with integers, $\mathbb{Z} = \{\ldots, -2, -1, 0, 1, 2, \ldots\}$. As in Chapter 3, divisors in divisibility statements are nonzero. The integer being divided may be zero.

❄ Definition 8.1 — Divisibility

For $d, a \in \mathbb{Z}$ with $d \neq 0$, write $d \mid a$ when

$$a = dk \quad \text{for some } k \in \mathbb{Z}.$$

We say that d **divides** a, that d is a **divisor** of a, or that a is a **multiple** of d. Write $d \nmid a$ when no such integer k exists.

The symbol $\mid$ states a relation between two integers; it does not denote the numerical quotient. The order matters: the number on the left is the divisor.

Example 8.1. We have $4 \mid 28$ because $28 = 4(7)$. The witness is 7. However, $4 \nmid 30$: the equation $30 = 4k$ forces $k = 15/2$, which is not an integer. Writing $30 = 4(15/2)$ is an equality of numbers, but it does not establish integer divisibility.

Also, $28 \nmid 4$. The claim $4 \mid 28$ cannot be reversed just by reversing the notation.

Example 8.2. Since $-24 = 6(-4)$, we have $6 \mid -24$. Since $24 = (-6)(-4)$, we also have $-6 \mid 24$. Neither the divisor nor its integer multiplier has to be positive.

For every nonzero integer d, $0 = d(0)$ gives $d \mid 0$. In particular, $1 \mid 0$, just as 1 divides every other integer. We do not use zero as a divisor.

Example 8.3. Let $d \neq 0$, and suppose $d \mid a$ and $d \mid b$. Prove that $d \mid (a - 2b)$.

Proof. There are integers j, k such that $a = dj$ and $b = dk$. Consequently,

$$a - 2b = dj - 2dk = d(j - 2k).$$

The multiplier $j - 2k$ is an integer. This is the required divisibility witness. □

The argument applies to all integers in the hypotheses. It also explains a useful general rule: integer combinations of multiples of d remain multiples of d.

8.2 Quotients and remainders

Exact divisibility does not always hold, but we can separate an integer into a multiple of a positive divisor and a bounded remainder. We use the following elementary result, often called the **division algorithm**.

❄ Definition 8.2 — Quotient and remainder

For each integer a and each positive integer n, there are unique integers q, r such that

$$a = nq + r, \qquad 0 \le r < n.$$

The integer q is the **quotient** and r the **remainder** when a is divided by n. We write $a \bmod n = r$ for this chosen remainder.

The bounds on r are part of the description. They choose the multiple nq at or immediately below a, so that $nq \le a < n(q+1)$. The distance from nq to a is the remainder. For a fixed positive n, $n \mid a$ exactly when that remainder is zero.

Example 8.4. A batch contains 47 records, grouped in packets of 6. Since

$$47 = 6(7) + 5, \qquad 0 \le 5 < 6,$$

there are seven complete packets and five records left. Thus $47 \bmod 6 = 5$ and $6 \nmid 47$. For comparison, $48 = 6(8) + 0$, so $48 \bmod 6 = 0$ and $6 \mid 48$.

Example 8.5. To divide -13 by 5, locate it between the consecutive multiples -15 and -10:

$$-13 = 5(-3) + 2, \qquad 0 \le 2 < 5.$$

The quotient is -3 and the remainder is 2.



The equality $-13 = 5(-2) - 3$ is also correct, but its final summand -3 is not the chosen remainder: it violates $0 \le r < 5$. We move to the preceding multiple of 5 and add 2.

Python (non-examinable)

For integer inputs and a positive divisor, Python's `divmod` returns the quotient and remainder with the bounds used here. The assignment `q, r = divmod(a, n)` stores the first returned value in `q` and the second in `r`.

```
a = -13
n = 5
q, r = divmod(a, n)
print(q, r)
print(n * q + r)
```

Output.

```
-3 2
-13
```

Thus $q = -3$ and $r = 2$, and the second line reconstructs $5(-3) + 2 = -13$. The positive-divisor condition matters: it ensures that the remainder lies between 0 and $n - 1$.

Example 8.6. Bob writes $29 = 6(3) + 11$ and calls 11 the remainder. The equality holds, but $11 \geq 6$. Move one further group of six into the quotient:

$$29 = 6(4) + 5.$$

Now $0 \leq 5 < 6$. Similarly, $-12 = 6(-2) + 0$ already has a valid remainder: a negative input can be an exact multiple.

Uniqueness of the remainder. Suppose $a = nq + r = nq' + r'$ with both remainders between 0 and $n - 1$. Then $n(q - q') = r' - r$. The right-hand side lies strictly between $-n$ and n. The only multiple of the positive integer n in that interval is 0. Thus $r' = r$, and then $q' = q$. Without the bounds, many decompositions would be possible.

8.3 Congruence

From now on our modulus n is an integer with $n \geq 2$. The quotient-and-remainder result also allows $n = 1$, but our modular systems will have at least two labels.

❄ Definition 8.3 — Congruence modulo n

Integers a, b are **congruent modulo** n if they have the same remainder on division by n. Write

$$a \equiv b \pmod{n}.$$

If their remainders differ, write $a \not\equiv b \pmod{n}$. The positive integer n is the **modulus**.

Example 8.7. We have

$$17 = 5(3) + 2, \qquad 7 = 5(1) + 2, \qquad -8 = 5(-2) + 2.$$

Therefore $17 \equiv 7 \pmod 5$ and $-8 \equiv 7 \pmod 5$. The three integers are distinct; their remainders on division by 5 are equal. In particular, the congruence $17 \equiv 7 \pmod 5$ does not assert $17 = 7$.

Example 8.8. For integers a, b and fixed $n \geq 2$, we claim

$$a \equiv b \pmod n \iff n \mid (a - b).$$

Proof. First suppose the remainders agree. Write $a = nq + r$ and $b = nt + r$. Then $a - b = n(q - t)$, an integer multiple of n.

Conversely, suppose $a - b = nk$ for some integer k. Write $b = nt + r$ with $0 \leq r < n$. Then

$$a = b + nk = n(t + k) + r.$$

This is a valid quotient-and-remainder description of a with the same remainder r. Uniqueness of the remainder gives $a \equiv b \pmod n$. □

Equivalently, $a \equiv b \pmod n$ means $a = b + nk$ for some $k \in \mathbb{Z}$. This form turns a congruence into an ordinary integer equation with a witness.

Example 8.9. We have $23 \equiv -1 \pmod 6$, because $23 - (-1) = 24 = 6(4)$. Nevertheless, $23 \bmod 6 = 5$, not -1. The chosen remainder must lie in $\{0, 1, 2, 3, 4, 5\}$. Both 23 and -1 have that same remainder 5.

Thus $\equiv$ compares two integers, whereas $a \bmod n$ names one particular integer in $\{0, \dots, n - 1\}$.

Example 8.10. The difference $17 - 7 = 10$ is divisible by 5, so $17 \equiv 7 \pmod 5$. It is not divisible by 4, so $17 \not\equiv 7 \pmod 4$. Stating that two integers are congruent without specifying a modulus leaves the relation unspecified.

8.4 Residue classes

Congruence generalises Chapter 7's even-difference relation. Modulo 2, equal remainders mean equal parity. For any fixed modulus, the same proof pattern establishes an equivalence relation.

Example 8.11. Fix $n \geq 2$ and let the domain be $\mathbb{Z}$. We use the divisibility test in Example 8.8.

Reflexivity. For every a, $a - a = 0 = n(0)$, so $a \equiv a \pmod{n}$.

Symmetry. If $a \equiv b \pmod{n}$, write $a - b = nk$ with $k \in \mathbb{Z}$. Then $b - a = n(-k)$, so $b \equiv a \pmod{n}$.

Transitivity. If $a \equiv b \pmod{n}$ and $b \equiv c \pmod{n}$, write $a - b = nj$ and $b - c = nk$ for integers j, k. Adding gives

$$a - c = (a - b) + (b - c) = n(j + k),$$

so $a \equiv c \pmod{n}$. All three properties hold on the whole domain.

❄ Definition 8.4 — Residue class

For $a \in \mathbb{Z}$, its **residue class modulo** n is

$$[a]_n = \{x \in \mathbb{Z} : x \equiv a \pmod{n}\} = \{a + nk : k \in \mathbb{Z}\}.$$

It is also called a **remainder class**. The integer a is a **representative** of the class.

The symbol $[a]_n$ denotes a set of integers. It is not the same object as the integer a. The subscript records which equivalence relation is being used.

Example 8.12. The classes modulo 4 are:

Chosen representative	Residue class
0	$[0]_4 = \{\ldots, -8, -4, 0, 4, 8, \ldots\}$
1	$[1]_4 = \{\ldots, -7, -3, 1, 5, 9, \ldots\}$
2	$[2]_4 = \{\ldots, -6, -2, 2, 6, 10, \ldots\}$
3	$[3]_4 = \{\ldots, -5, -1, 3, 7, 11, \ldots\}$

Every integer has exactly one remainder in $\{0, 1, 2, 3\}$. Consequently, every integer belongs to one of these classes and cannot belong to two different ones. The classes are nonempty, disjoint and together cover $\mathbb{Z}$: they form a partition, in Chapter 7's terminology. Each class is infinite, although there are only four classes.

Example 8.13. Since $-1 \equiv 3 \equiv 7 \pmod{4}$, we have

$$[-1]_4 = [3]_4 = [7]_4.$$

For instance, an integer differs from 7 by a multiple of 4 exactly when it differs from 3 by a multiple of 4: the two differences themselves differ by 4. The integers $-1, 3, 7$ are different representatives of one class; they do not name three new classes.

In general, there are exactly n distinct classes modulo n, represented once each by $0, \dots, n-1$. We write

$$\mathbb{Z}_n = \{[0]_n, [1]_n, \dots, [n-1]_n\}$$

for this set of classes. To keep computations short, we will also use the unique representatives as labels.

8.5 Arithmetic with congruences

Suppose $a \equiv b \pmod{n}$ and $c \equiv d \pmod{n}$, with the same modulus. Then

$$\begin{aligned} a + c &\equiv b + d \pmod{n}, \\ a - c &\equiv b - d \pmod{n}, \\ ac &\equiv bd \pmod{n}. \end{aligned}$$

These rules allow us to replace an integer by a congruent one inside a sum, difference or product.

Write $a - b = nj$ and $c - d = nk$ for integers j, k. The differences of the respective sums and differences are $n(j+k)$ and $n(j-k)$, hence are multiples of n. For products,

$$ac - bd = (a-b)c + b(c-d) = n(jc + bk).$$

Again the multiplier is an integer. The divisibility test therefore gives each congruence. Repeated multiplication also permits positive integer powers: if $a \equiv b \pmod{n}$, then $a^m \equiv b^m \pmod{n}$ for every integer $m \geq 1$.

Example 8.14. Modulo 5, we have $38 \equiv 3$ and $27 \equiv 2$. Therefore

$$38 + 27 \equiv 3 + 2 = 5 \equiv 0 \pmod{5},$$

and

$$38(27) \equiv 3(2) = 6 \equiv 1 \pmod{5}.$$

The sum has remainder 0 and the product has remainder 1. The ordinary product is 1026, but computing that full product was unnecessary for the remainder.

Example 8.15. Using the same inputs modulo 5,

$$38 - 27 \equiv 3 - 2 = 1 \pmod{5}.$$

In the reverse order,

$$27 - 38 \equiv 2 - 3 = -1 \equiv 4 \pmod{5}.$$

The last step uses $-1 = 5(-1) + 4$. Leaving the answer as a congruence to -1 is valid; the remainder itself is 4.

Example 8.16. Since $3^2 = 9 \equiv -1 \pmod{5}$,

$$3^8 = (3^2)^4 \equiv (-1)^4 = 1 \pmod{5}.$$

Thus 3^8 has remainder 1 when divided by 5. We used a small congruent value for 3^2 and then multiplied it four times. There was no need to expand 3^8 first.

Example 8.17. We have $2(1) \equiv 2(4) \pmod 6$, since $2 - 8 = -6$. Cancelling the factor 2 would give $1 \equiv 4 \pmod 6$, which is false: $1 - 4 = -3$ is not divisible by 6.

The multiplication rule proves a congruence after multiplying; it does not supply the converse. Indeed, from $6 \mid 2(1-4)$ we cannot conclude $6 \mid (1-4)$. Ordinary division by a nonzero number does not automatically preserve a congruence with the same modulus.

Example 8.18. Every integer x is congruent to one of $0, 1, 2, 3$ modulo 4. Squaring its representative gives:

$x \bmod 4$	0	1	2	3
$x^2 \bmod 4$	0	1	0	1

For example, $3^2 = 9 \equiv 1 \pmod 4$. The power rule ensures that every integer with representative 3 has its square in class $[1]_4$. The other columns work the same way. Thus every integer square is congruent to 0 or 1 modulo 4.

This is a proof for all integers, rather than a test of just four integers: every possible remainder is covered, and congruence preserves squaring. In particular, an equation $x^2 = 4k + 2$ has no solution with $x, k \in \mathbb{Z}$, since its right-hand side has remainder 2.

8.6 Modular systems

So far we have used congruence to compare integers. We can also calculate entirely with the finite set of representative labels. To describe this precisely, we must state the set and the operations used on it.

❄ Definition 8.5 — Binary operation and closure

A **binary operation on a set** S assigns exactly one element of S to each ordered pair of elements of S. In function notation it is a map $S \times S \to S$. For a proposed operation, **closure** means that every permitted pair has its result in S.

The word binary refers to two inputs. A **mathematical system** here means a specified set together with specified operations on that set. The set alone does not tell us which calculations are intended.

Example 8.19. Let $S = \{0, 1, 2, 3\}$. Ordinary addition gives $2 + 3 = 5 \notin S$. It therefore does not define a binary operation on S. Ordinary multiplication also fails closure, since $2(3) = 6 \notin S$. The fact that some pairs, such as $0 + 1$, stay in S does not establish closure for all pairs.

❄ Definition 8.6 — Modular addition and multiplication

Fix $n \geq 2$ and let $S_n = \{0, 1, \dots, n-1\}$. For $a, b \in S_n$, define

$$a \oplus_n b = (a + b) \bmod n, \qquad a \otimes_n b = (ab) \bmod n.$$


The symbols $\oplus_n$ and $\otimes_n$ denote addition and multiplication followed by reduction to the chosen remainder. Both operations have domain $S_n \times S_n$ and outputs in S_n.

Existence and uniqueness of a remainder ensure exactly one output in S_n for every input pair. Thus both are binary operations on S_n. The notation $(S_n, \oplus_n, \otimes_n)$ records the set and its two specified operations.

Example 8.20. In $S_4 = \{0, 1, 2, 3\}$,

$$2 \oplus_4 3 = 5 \bmod 4 = 1, \qquad 2 \otimes_4 3 = 6 \bmod 4 = 2.$$

The ordinary intermediate values 5 and 6 lie outside S_4; the operation outputs 1 and 2 lie inside it. Also, $3 \oplus_4 1 = 0$. This is an equality of representative labels under the defined operation; it does not assert the ordinary integer equality $3 + 1 = 0$.



Ordinary addition takes 3 to 4; adding 1 modulo 4 takes label 3 back to 0.

Operations on residue classes. The label $a \in S_n$ represents the class $[a]_n \in \mathbb{Z}_n$. On classes, adding $[a]_n$ to $[b]_n$ means forming $[a + b]_n$; multiplying means forming $[ab]_n$. These results do not depend on which integers represent the two classes.

For example, modulo 4, $[2]_4 = [6]_4$. Adding $[3]_4$ using the first representative gives $[2 + 3]_4 = [5]_4 = [1]_4$; using 6 gives $[6 + 3]_4 = [9]_4 = [1]_4$ again.

If $a \equiv a' \pmod{n}$ and $b \equiv b' \pmod{n}$, the arithmetic rules give

$$a + b \equiv a' + b' \pmod{n}, \qquad ab \equiv a'b' \pmod{n}.$$

So the resulting classes agree. Our operations on S_n record each resulting class by its unique label in $\{0, \dots, n-1\}$. We keep brackets for classes and use unbracketed labels when calculating with $\oplus_n, \otimes_n$.

Example 8.21. Here are both operations on S_4. The row gives the first input and the column gives the second.

$\oplus_4$	0	1	2	3
0	0	1	2	3
1	1	2	3	0
2	2	3	0	1
3	3	0	1	2

$\otimes_4$	0	1	2	3
0	0	0	0	0
1	0	1	2	3
2	0	2	0	2
3	0	3	2	1

For instance, row 3, column 2 in the addition table gives $3 \oplus_4 2 = 1$. Row 2, column 2 in the multiplication table gives $2 \otimes_4 2 = 0$, because $4 \bmod 4 = 0$. Two nonzero labels can therefore have product zero.

Each table covers all $4^2 = 16$ ordered input pairs, and every output lies in S_4. This is a complete finite check of closure. The general argument using bounded remainders already establishes closure for every S_n.

Python (non-examinable)

The symbol `%` computes a remainder in Python. With the positive modulus 4, it returns one of the labels $0, 1, 2, 3$. The outer loop chooses a row label a; the inner loop visits each column label b. Each empty list `row` receives the four outputs of addition modulo 4, using `append` to add each entry. Each `range(n)` supplies $0, 1, \ldots, n-1$, so here both loops use the labels $0, 1, 2, 3$.

```
n = 4
for a in range(n):
    row = []
    for b in range(n):
        row.append((a + b) % n)
    print(row)
```

Output.

```
[0, 1, 2, 3]
[1, 2, 3, 0]
[2, 3, 0, 1]
[3, 0, 1, 2]
```

These are exactly the rows of the addition table above. Replacing `a + b` by `a * b` generates the multiplication table on the same representative labels. The loops cover every ordered input pair in $S_4 \times S_4$.

❄ Definition 8.7 — Identity element

For a binary operation $\star$ on S, an **identity element** is an element $e \in S$ such that

$$e \star a = a \quad \text{and} \quad a \star e = a \qquad \text{for every } a \in S.$$

It leaves every element unchanged in either input position.

An identity belongs to a specified operation. There need not be one element that acts as the identity for every operation in a system.

Example 8.22. For $a \in S_n$, the integer a already satisfies $0 \le a < n$, so $a \bmod n = a$. Consequently,

$$0 \oplus_n a = a \oplus_n 0 = a, \qquad 1 \otimes_n a = a \otimes_n 1 = a.$$

Thus 0 is the additive identity and 1 the multiplicative identity for every $n \ge 2$. In the tables of Example 8.21, row and column 0 of the addition table reproduce their input labels. Row and column 1 of the multiplication table do likewise.

Zero is not a multiplicative identity: $0 \otimes_n 1 = 0 \neq 1$. One is not an additive identity: $1 \oplus_n 0 = 1 \neq 0$. Checking just an input that happens to stay unchanged would not establish the identity condition.

Example 8.23. On S_4, define $a \ominus_4 b = (a - b) \bmod 4$. This is a binary operation because its reduced output always lies in S_4. For every $a \in S_4$, $a \ominus_4 0 = a$. But $0 \ominus_4 1 = (-1) \bmod 4 = 3$, not 1. So 0 works on the right but is not a two-sided identity.

No other label can be a two-sided identity either. If e were such an identity, $e \ominus_4 0 = 0$ would force $e = 0$, which we have just ruled out. A set with a binary operation need not have an identity.

Example 8.24. Use addition modulo 4, but restrict the allowed labels to $A = \{1, 2, 3\}$. Since $1 \oplus_4 3 = 0 \notin A$, the restricted calculation is not a binary operation on A. It remains an operation on S_4. Specifying the modulus without the permitted input and output set would miss this change.

8.7 Exercises

8.1. Decide each divisibility claim, using the nonzero-divisor convention. Justify each answer.

a) $7 \mid -35$.

b) $-4 \mid 28$.

c) $9 \mid 0$.

d) $8 \mid 30$.

8.2. a) For integers d, a, b with $d \neq 0$, prove that $d \mid a$ and $d \mid b$ imply $d \mid (2a - 3b)$.

b) For nonzero integers d, e and an integer a, prove that $d \mid e$ and $e \mid a$ imply $d \mid a$.

8.3. For each pair (a, n), find q, r with $a = nq + r$ and $0 \leq r < n$.

a) $(53, 7)$.

b) $(-53, 7)$.

c) $(-42, 7)$.

d) $(0, 6)$.

e) $(18, 1)$.

8.4. The equalities below hold, but the final summand does not satisfy the required remainder bounds. Give a valid quotient and remainder for each.

a) $17 = 5(2) + 7$.

b) $-17 = 5(-3) - 2$.

c) $28 = 7(3) + 7$.

Explain why the bound is $r < n$, rather than $r \leq n$.

8.5. There are 1028 records, packed into full groups of 32. Find the number of full groups and the number left over. State whether $32 \mid 1028$ and justify the conclusion from your quotient-and-remainder equation.

8.6. State TRUE or FALSE for each claim, justifying it by divisibility of the difference.

a) $14 \equiv -1 \pmod 5$.

b) $18 \equiv 2 \pmod 4$.

c) $-12 \equiv 3 \pmod 5$.

d) $7 \equiv 0 \pmod 3$.

e) $0 \equiv 6 \pmod 6$.

f) $11 \equiv 5 \pmod 4$.

8.7. Show that $20 \equiv -1 \pmod 7$. Find $20 \bmod 7$ and $(-1) \bmod 7$. Why is -1 a valid congruent integer but not a valid chosen remainder modulo 7?

8.8. Show that $5 \equiv 8 \pmod 3$ but $5 \not\equiv 8 \pmod 6$. Explain why these conclusions are compatible. Give another pair of distinct integers with the same two properties.

8.9. Let $n \geq 2$, and suppose $a \equiv b \pmod n$ and $b \equiv c \pmod n$. Use equations $a - b = nj$ and $b - c = nk$ to prove $a \equiv c \pmod n$. Identify which relation property this establishes and why $j + k$ is an allowed witness.

8.10. List the three distinct classes modulo 3, giving a set-builder description of each. For each class, include two negative and two nonnegative members. Explain why every integer belongs to exactly one class.

8.11. Decide whether each equality is correct and justify it. For a failed equality, give an integer that belongs to one set but not the other.

a) $[2]_5 = [12]_5$.

b) $[-1]_5 = [4]_5$.

c) $[0]_4 = [2]_4$.

d) $[1]_3 = [1]_5$.

8.12. List all integers x with $-15 \le x \le 15$ such that $x \equiv 2 \pmod 5$. Give an equation $x = 2 + 5k$ and identify all integer values of k that produce your list.

8.13. Find the chosen remainder in each case, showing congruence steps.

a) $(46 + 39) \bmod 7$.

b) $(46 \cdot 39) \bmod 7$.

c) $(39 - 46) \bmod 7$.

d) $(4 - 19) \bmod 6$.

8.14. Suppose $a \equiv 2 \pmod 5$ and $b \equiv 4 \pmod 5$. Find the chosen remainders of $a + b$, ab and $a^2 + b^2$ modulo 5. Explain why your answers do not depend on the particular integers represented by a and b.

8.15. Show that $2^3 \equiv 1 \pmod 7$. Use $20 = 3(6) + 2$ to find $2^{20} \bmod 7$, without expanding 2^{20}. State where repeated multiplication of congruences is used.

8.16. Suppose $a - b = nj$ and $c - d = nk$, with all letters integer and $n \ge 2$. Express $ac - bd$ as n times an integer to prove that $a \equiv b \pmod n$ and $c \equiv d \pmod n$ imply $ac \equiv bd \pmod n$. Explain why this also makes multiplication of residue classes independent of representatives.

8.17. Check that $4(2) \equiv 4(5) \pmod 6$. Charlie cancels the nonzero factor 4 and concludes $2 \equiv 5 \pmod 6$. Show that this conclusion is false. Explain which implication the valid multiplication rule supplies and which Charlie has assumed.

8.18. On $S_5 = \{0, 1, 2, 3, 4\}$, which of the following rules define binary operations? For a failed rule give an input pair whose result lies outside S_5; for a successful rule justify all input pairs.

a) Ordinary addition $a + b$.

b) Ordinary multiplication ab.

c) $(a + b) \bmod 5$.

d) $(ab) \bmod 5$.

8.19. Construct the addition and multiplication tables on $S_3 = \{0, 1, 2\}$ using $\oplus_3$ and $\otimes_3$. Explain how the entries show closure. Use the tables to identify the additive and multiplicative identities, checking both the relevant row and the relevant column.

8.20. In S_5, Eve notes that $2 \otimes_5 0 = 0$ and claims that 2 is a multiplicative identity. Give an input that disproves her claim. State the full identity condition and identify the actual multiplicative identity.

8.21. On S_5, define $a \ominus_5 b = (a - b) \bmod 5$.

a) Calculate $1 \ominus_5 4$ and $4 \ominus_5 1$.

b) Explain why the operation is closed.

c) Show that $a \ominus_5 0 = a$ for every $a \in S_5$, but 0 is not a two-sided identity.

d) Explain why no other element can be a two-sided identity.

8.22. Use addition and multiplication modulo 4.

a) On $A = \{1, 2, 3\}$, give a pair showing that addition is not closed.

b) On $B = \{0, 2\}$, check every ordered pair to decide whether addition is closed.

c) On the same B, check every ordered pair to decide whether multiplication is closed.

8.23. On $S = \{u, v\}$, define $\star$ by the following table, with the first input selecting the row.

$\star$	u	v
u	u	v
v	v	u

Explain why $\star$ is a binary operation on S. Identify its identity and show why the other element is not an identity. The labels u, v are elements of this set; no ordinary numerical addition is being assumed.

8.24. The printer-checking order is Alice, Bob, Charlie, David, repeating, with labels $0, 1, 2, 3$ respectively. At turn 0, Charlie checks.

a) Give a formula for the checking label at turn $t \geq 0$ using a remainder.

b) Who checks at turn 14?

c) If the same repeating allocation is extended backwards to integer turns, who checks at turn -3?

Explain why including the initial label in the formula is necessary.

8.25. For any integer k, its remainder modulo 2 is 0 or 1. Use these two cases and congruence arithmetic to prove $2 \mid k(k+1)$ for every $k \in \mathbb{Z}$. Explain why two remainder cases cover infinitely many integers.

8.26. Use Example 8.18 to find all possible remainders of $x^2 + y^2$ modulo 4, where $x, y \in \mathbb{Z}$. Prove that there are no integers x, y, m satisfying

$$x^2 + y^2 = 4m + 3.$$

Explain why this is a general impossibility proof rather than a check of a few values of x and y.